




Oxford University Hospitals
NHS Foundation Trust

Azithromycin and Lung Disease

Information for patients



What is azithromycin?

Azithromycin is a type of antibiotic called a 'macrolide antibiotic'. Antibiotics are used to treat infections caused by microorganisms, such as bacteria.

Why have I been prescribed azithromycin?

You have been prescribed azithromycin as a preventative treatment, to try to improve control of your chest symptoms and exacerbations (lung attacks).

How does it work?

Azithromycin kills certain bacteria and reduces inflammation in the lungs, which may help to reduce the number of lung attacks you have. Azithromycin may help reduce chest symptoms, such as coughing, sputum (phlegm) production and breathlessness.

What do I need to know before I take azithromycin?

Do not take this medication if you are allergic to azithromycin, any of its ingredients or any other macrolide antibiotic, such as erythromycin or clarithromycin.

Your doctor will usually ask you to provide a sputum sample, to check for non-tuberculosis mycobacterium, which is a similar bacteria related to tuberculosis. It is important that we check for this prior to starting azithromycin, as if these bacteria are present azithromycin can't be used.

The doctors will also carry out a heart trace (ECG) before starting this treatment, to check the rhythm of your heart. You will also have blood tests to check your liver function, so that we can see if the azithromycin starts to affect the organ. These tests will be repeated at the hospital after you have been taking the azithromycin for 1 month to ensure that the rhythm of your heart and your liver function have not been affected by the azithromycin.

Further monitoring will be carried out by your GP as your treatment with azithromycin progresses.

What dose will I need to take?

The usual dose is one 250mg capsule, three times a week (usually Monday, Wednesday and Friday), but you may be prescribed a higher dose (one 500mg capsule, three times a week).

Are there any side effects?

Azithromycin doesn't usually cause any side effects.

Less than 10 in 100 people experience tummy upset, diarrhoea, constipation, dizziness, headaches or a skin rash.

Less than 1 in 100 people report some hearing loss. The first symptom of this can be tinnitus (ringing in the ears). If you experience this symptom the azithromycin should be stopped immediately. Your hearing will then gradually return to normal.

Less than 1 in 10,000 people experience changes in their heart tracing or an effect on their liver. Your doctor is likely to repeat your heart tracing and blood tests once you have started azithromycin treatment, to check on this.

For further information on side effects please see the manufacturer's patient information leaflet, which will be in the box of medication.

What if I forget to take my dose?

This should not cause any problems. Just take your next dose as scheduled.

How long will I be on this medication?

You are likely to need to take azithromycin for 3 to 6 months (if it isn't causing any significant side effects). We will then review whether it is working. If it is reducing the number of lung attacks that you suffer from, and reducing your symptoms (coughing, sputum production and breathlessness), then you can continue taking it.

Can azithromycin affect any of my other medications?

Please let your hospital doctor and GP know all the medications you are taking before you start azithromycin, including over the counter and herbal remedies. They will need to check for any potential interactions (reactions between the medications and azithromycin). Always speak to your doctor and/or pharmacist before starting any new medicine.

If you are taking any of the following medications you will need extra blood tests, to check that the other drugs are working and not causing side effects. The doses of these medicines may also need to be adjusted.

Theophylline and aminophylline

Azithromycin can increase the levels of these medicines in your blood, so you may need extra monitoring with blood tests. The brand name of this medicine is Uniphyllin.

Warfarin

Azithromycin can increase the blood levels of this medication and extra monitoring with blood tests may be required, especially when azithromycin is first started.

Digoxin

Azithromycin can increase the blood levels of this medication and extra monitoring may be required, especially when azithromycin is first started.

Statins

Azithromycin does not increase the blood level of statins in the blood, but you should look out for signs/symptoms of muscle aches, pains or weakness (rhabdomyolysis) and report any of these symptoms to your GP if they occur.

What should I do if I suffer a flare-up of my lung disease?

Although azithromycin is an antibiotic, you will still need to take other antibiotics for any chest infections. You should stop taking your azithromycin whilst on the other antibiotics. You can then start to take it again when they are finished.

If I need surgery should I stop taking azithromycin?

You should not have to stop taking this medication before any surgery you might need. If you are unsure, check with your respiratory consultant or the surgical team.

Can azithromycin affect fertility or pregnancy?

Azithromycin has no known effect on fertility. However, due to a lack of clinical data, if you become pregnant whilst you are taking azithromycin, stop taking it and tell your GP.

Can I drink alcohol whilst taking azithromycin?

Yes, alcohol doesn't affect azithromycin. However, it may affect other antibiotics or medications you might be taking. Speak with your doctor or GP if you are unsure.

Further questions

If you have any other questions, please speak to the clinician who prescribed you the azithromycin.

Further information

If you would like an interpreter, please speak to the department where you are being seen.

Please also tell them if you would like this information in another format, such as:

- Easy Read
- large print
- braille
- audio
- electronic
- another language.

We have tried to make the information in this leaflet meet your needs. If it does not meet your individual needs or situation, please speak to your healthcare team. They are happy to help.

Author: Respiratory Medical Team
January 2024
Review: January 2027
Oxford University Hospitals NHS Foundation Trust
www.ouh.nhs.uk/information



Making a difference across our hospitals

charity@ouh.nhs.uk | 01865 743 444 | hospitalcharity.co.uk

OXFORD HOSPITALS CHARITY (REGISTERED CHARITY NUMBER 1175809)

