



Oxford University Hospitals
NHS Foundation Trust

Discharge advice after your ERCP

Information for Patients



We will discuss your ERCP result with you before you leave and give you a copy of your report. Please ask if you have any questions.

Your ERCP

You have had an ERCP:

under midazolam and fentanyl sedation

under propofol sedation

The sedation you have been given makes you sleepy and can cause forgetfulness and lack of concentration for up to 24 hours. You should go home and rest today.

You must be collected and accompanied home.

If you have had sedation and live alone, we recommend that you try to arrange for someone to stay with you overnight.

For 24 hours after the sedation you must not:

- drive
- drink alcohol
- operate heavy machinery
- sign any legally binding documents.

Most people feel back to normal after 24 hours.

After effects

You may have a sore throat for about 24 hours. This is quite normal.

During the ERCP you had a:

Sphincterotomy

Pre-cut

Stent

We will explain to you what this procedure involved and any treatment you may need afterwards.

Please contact the Urgent Endoscopy Advice Line on the number at the back of this leaflet if you have:

- pain in your neck, chest or abdomen, especially epigastric pain (pain in the centre of your chest/abdomen) or pain that radiates to your back
- nausea or vomiting
- a fever (hot or cold spells)
- any signs of bleeding, including black faeces or dizziness.

What happens next?

A copy of the endoscopy report will be sent to your GP electronically within 1 working day. You should also be given a copy of the report before you go home.

Your results

If the investigation was requested by a hospital Consultant, the report will be sent to them. The Consultant will then write to you and your GP to tell them the results of the biopsy. If you are having other tests your Consultant may need to wait for all the results before they write to your GP. This may not be until after you have returned for a further outpatient appointment.

Your follow up is:

- No further appointment is needed
- An outpatient appointment will be sent to you in weeks/months
- A repeat procedure will be arranged in weeks/years
- The following further investigations will be arranged:

.....

.....

.....

How to contact us

If you have any pain in your neck chest or abdomen, vomiting, fever or signs of bleeding, please contact the Urgent Endoscopy Advice Line (24 hour).

Tel: **01865 741 166** – ask to bleep **6825**

Let us know your views and experience

You can comment by visiting: **ouh.nhs.uk/fft**

Or you can email: **endoscopy@ouh.nhs.uk**

Or write to:

Endoscopy Sister
Endoscopy Department
Level 2
John Radcliffe Hospital
Headley Way
Oxford OX3 9DU

Further Information

Please speak to the department where you are being seen if you would like an interpreter. You will find their contact details on your appointment letter. Please also ask them if you would like this information leaflet in another format, such as:

- easy read
- large print
- braille
- audio
- electronically
- in another language.

We have tried to make this information meet your needs. If it does not meet your individual needs or situation, please speak to your healthcare team. They will be happy to help.

Author: Endoscopy Team
April 2021
Review: April 2024
Oxford University Hospitals NHS Foundation Trust
www.ouh.nhs.uk/information



Making a difference across our hospitals

charity@ouh.nhs.uk | 01865 743 444 | hospitalcharity.co.uk

OXFORD HOSPITALS CHARITY (REGISTERED CHARITY NUMBER 1175809)

