


How to Recognise and Manage Allergic Reactions

Information for patients



What is an allergic reaction?

An allergic reaction occurs when the body's immune system reacts to a normally harmless substance. An allergic person's immune system considers these substances to be potentially damaging and releases histamine to defend the body against them. This causes certain changes in the body, which may produce a variety of mild to severe symptoms. This allergic response can develop after exposure by skin contact, swallowing or inhalation (breathing in).

Anaphylaxis or anaphylactic shock is a condition where the body's immune system severely over-reacts to something. This results in an acute, severe, potentially (but rarely) fatal allergic reaction, which needs immediate medical attention.

Causes of allergic reactions

Certain foods

- It is the protein in the food that is usually the allergic part.
- Common allergic foods are nuts (especially peanuts), fish, shellfish, eggs and cow's milk.
- Most allergic food reactions occur immediately after swallowing. Only rarely do reactions begin several hours after eating.
- Food allergies are more common in people, who have other allergic conditions such as asthma, eczema and hay fever.

Insect stings

- In particular wasp and bee stings.
- The reaction to an insect sting is usually immediate (within 30 minutes).

Natural rubber latex

- Some common sources of latex:
 - Contraceptives: condoms, diaphragms.
 - Balloons and rubber bands.
 - Carpet backing and furniture filling.
 - Rubber gloves.
 - Medical/dental items such as catheters, gloves, disposable items.

Medicines

- Anaphylactic reactions usually occur after the first dose of a course.

Idiopathic

- Occasionally, especially in adults, a cause cannot be identified.

Avoiding Allergens

The most important aspect of managing an allergy is to AVOID the substance that you are allergic to (the allergen). You will not experience an allergic reaction if you AVOID the allergen(s).

Food

When a food allergy has been identified, it is essential that you avoid eating even a tiny quantity of the food that you are allergic to (unless advised differently by the allergy clinic). An allergic reaction can also happen after skin contact with that particular food.

A major problem with foods is the accidental exposure to a particular food allergen (especially peanuts) is possible when it is a concealed or an undisclosed ingredient. It is important to:

- Read all labels carefully, even if the product has been eaten before, as recipes change.
- Be more careful when eating out, on holidays, on outings and at parties or functions. These are times when accidental ingestion can occur more easily. Be alert!
- Ask about ingredients in food when eating from restaurants, ice-cream parlours, in-store bakeries and delicatessen counters (foods are generally unlabelled in these places). Stress the seriousness of the allergic condition to the staff. Preferably talk directly with the chef if necessary – ask about the ingredients and stress the use of clean utensils. If they can not guarantee that a specific dish is safe, it is best to eat elsewhere.
- Ask retailers and manufacturers for lists of produce, which are free from certain ingredients, such as peanuts, nuts, egg, and milk.
- Dieticians are able to give advice about avoidance of particular foods and have very useful information leaflets.

- Cross contamination is a risk for individuals with food allergy. This happens when a food has unknowingly come in contact with the allergenic food. Ideas to reduce the risk of cross contamination include:
 1. Take extra care when preparing food so that contaminated cutlery, crockery, or work surfaces do not come into contact with the 'safe' food. For example, use clean utensils and wipe down surfaces with hot soapy water.
 2. Keep any allergenic food separate from other foods.
 3. If a partner, relative or friend has been eating the allergenic food, ask them to wash their hands and rinse their mouth well before touching or kissing.

Insect stings

Here are some steps that may help prevent you being stung (by wasps in particular):

- Avoid dressing in brightly coloured clothing in the warmer months.
- Ensure you wear shoes when outside.
- Avoid wearing perfumes and fragrances.
- Avoid eating food (especially sweet ones) outside.
- Avoid canned drinks in the wasp season; boxed drinks with straws may be safer.
- When outside avoid open rubbish bins and keep food covered.
- Avoid picking fruit including fallen fruit.
- Ensure nests are removed promptly.
- Keeping doors/windows closed in summer and/or use fly nets.
- Use insect repellents.

Natural rubber latex (see separate latex information sheet)

A severe allergic reaction is most likely to occur when latex has come into contact with mucous membranes or directly with tissue (as during surgery). It is important to:

- Always warn doctors, dentists, paramedic staff and hairdressers if you are allergic to latex.
- Be aware of all substances that may contain latex.

Medicines

- Ensure that you know all the names of the medicines that you are allergic to as medicines sometimes have more than one name.
- Always warn doctors, dentists, nurses and paramedic staff what you are allergic to. Wearing an emergency medical alert bracelet recording these is helpful.
- Always check any medicines prescribed to you by your GP or in hospital.

Signs and symptoms

Despite avoiding the allergic substances, accidents can happen. In an allergic reaction any of the following symptoms may occur:

Mild/moderate symptoms

- Tingling, itching or burning sensation in the mouth (this is a useful warning that you have eaten a food you are allergic to).
- A nasty taste in the mouth.
- Itching of skin, throat and/or eyes.
- Rapid development of nettle rash/wheals/hives (urticaria).
- Swelling, particularly of the face and lips.
- Feeling hot or chilled.
- Rising anxiety.
- Flushed.
- Tummy, abdominal or stomach pain.
- Nausea and/or vomiting.
- Mild wheeziness.

Severe symptoms (known as anaphylaxis)

The presence of any one or several of these features indicates an anaphylactic reaction, which needs to be treated immediately with Epinephrine (Adrenaline):

- Severe swelling of the tongue or throat associated with difficulty in breathing.
- Difficulty talking or hoarse voice.
- Severe wheeze or difficulty breathing.
- Faintness or dizziness.
- Looking very pale or blue and clammy.
- Disorientated or unresponsive.
- Collapse.

Milder reactions are much more common. Anaphylaxis, the most severe type of allergic reaction, is uncommon. It can be life threatening but is very rarely fatal.

Most reactions occur quickly after ingestion or contact with the allergen. Only rarely do they occur a few hours later. There can also be a second phase of the reaction when symptoms recur. This usually occurs within 6 hours of the onset of the reaction. This is called a late phase reaction.

Actions

Always have antihistamines, and if prescribed, a blue reliever inhaler and/or EpiPen available.

Actions for mild/moderate symptoms

1. Take an antihistamine such as Chlorphenamine (Piriton), Cetirizine (Zirtek) or Loratadine (Clarityn). (Cetirizine or Loratadine may be better, because they are non-sedating antihistamines and do not cause drowsiness.)
2. Treat mild wheeziness with a blue reliever inhaler (if you have one) such as Salbutamol (Ventolin) or Terbutaline (Bricanyl).
3. Let someone know that you are having a reaction and ask them to stay with you to help monitor the reaction to make sure it is getting better, not worse.

Actions for severe symptoms

Remember mild symptoms may develop into a more serious reaction.

1. If the symptoms are or do become severe, as listed earlier, treatment is with an injected drug called Adrenaline. The recommended way of giving adrenaline is with an EpiPen Autoinjector. This is a pre-filled syringe with a concealed spring-activated needle, which automatically injects the correct dose of Adrenaline when the device is pushed firmly against the thigh. The enclosed action plan gives more details on how to administer an EpiPen and a more detailed management plan.
2. Phone **999 and ask for an ambulance** with a paramedic crew stating that it is a severe allergic reaction.
3. Lying down may help if you feel dizzy or light-headed. If you are breathless, you may feel more comfortable sitting up.
4. Continue to monitor the condition. If there is no improvement after 10 minutes, a second dose of Adrenaline can be given via another EpiPen. (One dose per EpiPen). Note the time of both doses.

Things to remember about the EpiPen

When is Epinephrine (Adrenaline) given?

It should be given immediately to anyone having a severe attack. Do not be afraid to use your Epinephrine (Adrenaline). If someone is severely unwell with an allergic reaction, they are more at risk from the reaction than from the EpiPen. It is the most important treatment in stopping a severe reaction and should be given as soon as possible. You should have practiced with a dummy EpiPen, so that you know what to do.

Epinephrine (Adrenaline) may well make you feel more anxious, increase your heart rate, give you a few palpitations, make your breathing a bit more rapid and make you feel nauseated.

Patient training

We will show you how to use the EpiPen with a dummy version.

More information can be viewed at: www.EpiPen.co.uk

Dummy demonstrators can also be obtained through this website or at:

MEDA Skyway House,
Parsonage Road, Takeley
Bishop's Stortford, CM22 6PU
Telephone: +44 (0)845 460 0000

Carrying the EpiPen

You should carry your EpiPen at all times. It should remain in the protective yellow case as the Adrenaline is sensitive to light. It should also be kept at room temperature to prevent the contents from discolouring. It should never be stored in a fridge or left in a hot place.

Replacing the EpiPen

It is your responsibility to replace the EpiPen when it has expired, if the contents have discoloured or if the EpiPen has been used. Make a note on a calendar or in your diary to remind yourself.

Alternatively, the manufacturers of the EpiPen provide a service where you fill in your details with the expiry date of your EpiPen either on a card or online and they will send you a reminder just before it expires.

Travelling aboard

When going on holiday take 2 EpiPens with you. If travelling by air, make sure your antihistamines, inhalers and EpiPens are in your hand luggage. Some airlines may require a covering letter from your GP when carrying injections (or you can use a copy of your Allergy Clinic letter). Check with the booking agency.

If travelling aboard, find out the emergency telephone number for the specific country you are going to and where the nearest hospital is in relation to your holiday destination. It is always best to be prepared! Ensure your travel insurance is aware of your allergy.

Emergency medical alert bracelets

These can be obtained from many different companies, but these are a few we have found:

Medic Alert Foundation

Suite 1, Liscombe South

Liscombe Park

Soulbury, LU7 0JL

Telephone: 01908 951 045

Website: www.medicalert.org.uk

Medi-Tag

Medi-Tag

PO Box 17000

Birmingham, B32 9HR

Telephone: 0121 200 1616

Website: www.medi-tag.co.uk

Further information

Contact the Dermatology Department at the Churchill Hospital:

Churchill Hospital

Dermatology Department
Oxford, OX3 7LJ

Telephone: **01865 228 224**

Consultants:

Dr J Reed

Consultant Dermatologist with an interest in allergy.

Dr R Jain

Consultant Immunologist.

Dr S Patel

Consultant Immunologist.

Dr V Crump

Consultant Allergist.

Dr J. Miller

Consultant Immunologist.

The Anaphylaxis Campaign

Anaphylaxis UK
1 Alexandra Road
Farnborough
Hampshire, GU14 6BU

Telephone Helpline: 01252 542 029

Website: www.anaphylaxis.org.uk

Allergy UK

Allergy UK
London House Business Centre
Texcel Business Park
Thames Road
Crayford, DA1 4SL
Telephone Helpline: 01322 611 641
Website: www.allergyuk.org

Asthma & Lung UK

The White Chapel Building
10 Whitechapel High Street
London, E1 8QS
Telephone Helpline: 0300 222 5800
Website: www.asthmaandlung.org.uk

National Eczema Society

11 Murray Street
London, NW1 9RE
Telephone: 020 7281 3553
Helpline: 0800 448 0818
Website: www.eczema.org

BAD

British Association of Dermatologists
Willan House
4 Fitzroy Square
London, W1T 5HQ
Telephone: 020 7383 0266
Website: www.bad.org.uk

Further information

If you would like an interpreter, please speak to the department where you are being seen.

Please also tell them if you would like this information in another format, such as:

- Easy Read
- large print
- braille
- audio
- electronic
- another language.

We have tried to make the information in this leaflet meet your needs. If it does not meet your individual needs or situation, please speak to your healthcare team. They are happy to help.

Author: Dr. John Reed, Consultant Dermatologist with an interest in allergy.

July 2024

Review: July 2027

Oxford University Hospitals NHS Foundation Trust

www.ouh.nhs.uk/information



Making a difference across our hospitals

charity@ouh.nhs.uk | 01865 743 444 | hospitalcharity.co.uk

OXFORD HOSPITALS CHARITY (REGISTERED CHARITY NUMBER 1175809)

